

Interpretation of the Spatial Power Spectra of Neutral Hydrogen in the Galaxy and in the Small Magellanic Cloud

Itzhak Goldman

School of Physics and Astronomy, Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv 69978, Israel, email: goldman@wise1.tau.ac.il

ABSTRACT

Recent 21 cm radio observations of H_I regions in the Small Magellanic Cloud, have revealed spatial power spectra of the intensity, which are quite similar in shape to those previously deduced for the Galaxy. The similarity, in spite the differences in the physical parameters between the Galaxy and the SMC, suggests that the shape of the power spectra reflects some underlying mechanism which is not too sensitive to the environmental specifics. In this paper we present an interpretation for the observational power spectra in terms of a large scale turbulence in the interstellar medium, in which the emitting H_I regions are embedded. The turbulence gives rise to density fluctuations which lead to the observed intensity fluctuations, in the H_I regions. The observational power spectra are used to deduce the turbulence spectral function. In the SMC, the turbulence largest eddies are comparable in scale to the SMC itself. This implies that turbulent mixing should have smoothed out any large scale abundance gradients. Indeed, this seems to be the case, observationally.

The turbulence is also expected to amplify and shape up the large scale magnetic field. Indeed, the observational data indicate the existence of a large scale disordered field of the strength expected from energy equilibrium with the turbulent velocity field. The large scale turbulence is most probably generated by instabilities in the large scale flows induced by the tidal close encounter with the LMC $\sim 2 \times 10^8$ yr ago. The life-time of the largest eddies is $\sim 4 \times 10^8$ yr so the turbulence had not yet enough time to decay and persists even though the energy source is no longer there.

Subject headings: ISM: structure – turbulence – ISM: kinematics and dynamics – ISM: abundances – ISM: magnetic field

1. Introduction

Crovisier & Dickey (1983), and Green (1993) have used 21 cm radio observations to derive the intensity spatial power spectra for our Galaxy. Recently, Stanimirović *et al.* (1999) obtained H_I spatial power spectra for the Small Magellanic Cloud (SMC). Some of the observations were interferometric and yielded the power spectrum directly, and some single dish observations from which the power spectrum was calculated by Fourier transform.

The resulting power spectra exhibit a power-law behavior, $p(q) \propto q^\gamma$, over a wide range of spatial scales: ~ 10 – 200 pc in the Galaxy, and ~ 30 pc– 4 kpc for the SMC. Here q is the absolute value of the 2-dimensional wavenumber in the plane of the sky, and the observational power-law indices are $\gamma \sim -(2.8 \div 3)$ for the Galaxy and $\gamma \sim -3$ for the SMC.

Given the differences in the spatial scales for which the spectra were obtained, and the differences in the physical parameters between the Galaxy and the SMC, the similarity of the indices is remarkable. It suggests the existence of some underlying mechanism that is not sensitive to the specific physical parameters of the environment. Stanimirović *et al.* (1999) attributed the power-law shape to a fractal structure of H_I clouds hierarchy.

The observational power spectra reveal the existence of intensity correlations over a wide range of spatial scales, and this naturally suggests that turbulence in the interstellar medium (ISM) is the sought for underlying mechanism. Such an approach was suggested by Lazarian (1995) and followed up by Lazarian & Pogosyan (2000). In this paper we present an interpretation of the observational power spectra in terms of density fluctuations which are generated by a large scale turbulence in the ISM, in which the emitting neutral hydrogen regions are embedded. The model determines the power spectrum for any given spectral function of the underlying large scale turbulence. Therefore, the observational power spectra can be used to study the above large scale turbulence. From the observational power spectra, we deduce the spectral function of the underlying turbulence: $F(k) \propto k^{-2}$, with k denoting the absolute value of the 3-dimensional wavenumber. As detailed in the Discussion section, this spectral function is quite common in various components of the ISM and characterizes turbulence in a compressible medium.

2. The Model Outline

The observed lines of sight are generally optically thin in the 21 cm line. For the SMC, Stanimirović *et al.* (1999) note that self absorption effects are limited to narrow ranges in velocities and directions. They find that for most of the covered area the emission is from regions with column densities, integrated over the entire velocity span, of $N_{H_I} < 2.5 \times 10^{21} \text{ cm}^{-2}$. Each of the power spectra corresponds to a velocity range which equals 1/6 of the entire velocity span. Thus, one can estimate that typically each power spectrum corresponds to a column density $N_{H_I} \sim 4 \times 10^{20} \text{ cm}^{-2}$.

For the outer arms of the Galaxy, for which Green (1993) obtained the best fitting power spectra, we can estimate the column density using the relation (Spitzer 1978)

$$N_{H_I} = 1.82 \times 10^{18} \text{ cm}^{-2} \int T_B dv \quad (1)$$

where T_B and v are in K and km/s , respectively. For the outer arms, the average

$T_B \sim 15K$ over velocity groups of 3 channels, each of width $2.64km/s$ (Green 1993), yields $N_{H_I} \sim 2 \times 10^{20} cm^{-2}$.

To estimate the physical depth D of the emitting neutral hydrogen region, contributing to each power spectrum, we take a number density of neutral hydrogen of $n = 10cm^{-3}$. This value is intermediate between $n \sim 5cm^{-3}$ corresponding to warm neutral hydrogen with temperature $T \lesssim 200K$, and $n \gtrsim 30cm^{-3}$ of the cool neutral hydrogen with $T \lesssim 100K$ (Wolfire *et al.* 1995). This translates for the Galaxy outer spiral arms to a physical depth $D \sim 7pc$ and for the SMC to $D \sim 15pc$.

The power spectra were grouped by the observers in radial velocity intervals. It is interesting that, both for the Galaxy and the SMC, the power spectra for the different velocity groups are similar. In the Galaxy, adopting a rotation curve to deduce distances, each such velocity group represents a range of distances. Thus, in Green (1993) this range for the outermost spiral arms is $\simeq 2.2kpc$ while for the Perseus data the range is $\simeq 0.6kpc$. Each such distance range is *much larger* than the above estimate of $D \sim 7pc$. Therefore, it follows that the intensity is contributed by *disjoint subregions* of neutral hydrogen with total depth equaling D , with each subregion having a depth which is only a fraction of D , e. g. $\sim 1pc$.

In the SMC, the total velocity span probably represents large scale flows, of few kpc, that are the result of tidal interactions with the LMC and with the Galaxy. Indeed, N-body simulations by Gardiner & Noguch (1996) indicate a very strong tidal impact imparted to the SMC by the LMC about $\sim 2 \times 10^8 yr$ ago, with a duration of $\sim 1 \times 10^8$ yr. The resulting simulated velocity field exhibits velocities in the range of tens of km/s. This is consistent with the observational velocity span of $\sim 80km/s$ of the neutral hydrogen emission as reported by Stanimirović *et al.* (1999). The velocity range of each of the power spectra, corresponds to a distance interval $\sim 1kpc$ which is *much larger* than the total depth of the emitting regions, $D \sim 15pc$ contributing to each power spectrum. Therefore in this case too, the emission arises from a number of disjoint zones, each of depth which is a fraction of D , e. g. $\sim 2pc$.

We note that in both cases, the depth of each of the disjoint emitting regions is also *much smaller* than the scales, in the plane of the sky, probed in the power spectra: for the Galaxy up to ~ 200 pc and for the SMC up to 4 kpc. Therefore, the emitting H_I regions occupy a small fraction of the ISM volume, in which they are embedded. A small space filling factor of neutral hydrogen is implied also in the fractal interpretation of Stanimirović *et al.* (1999).

In the optically thin case, the intensity is proportional to the column density of neutral hydrogen in the line of sight (Spitzer 1978). Thus, the intensity fluctuations are due to density fluctuations in the disjoint neutral hydrogen regions. In the present model, the density fluctuations are regarded to be a response to a large scale 3-dimensional turbulence in the ISM, in which the neutral hydrogen regions are embedded.

Since the observational optical depths are small, the following picture suggests itself : the H_I subregions (clouds) are waved by a large scale turbulence in the embedding medium. In this picture the neutral hydrogen regions are regarded as “passive markers” which follow the turbulence field but do not feed back on it dynamically. The turbulence induces density fluctuations in the neutral hydrogen subregions that manifest as intensity fluctuations.

To be specific, we consider a simple model of density fluctuations in thin slabs or sheets parallel to the plane of the sky, which are separated along the line of sight direction by distances *much larger* than their depth. This geometry is consistent with the above estimates of the depth of the emitting subregions. H_I regions are generally assumed to have a filamentary structure and filaments are expected to have random orientations. This is not to say that all H_I emitting regions are necessarily filamentary. However, in our case the contributions to the power spectrum from a given angular separation in the plane of the sky will predominantly be those from regions at roughly the *same* distance along the line of sight. So, instead of summing over the actual filaments, contributing to each power spectrum, we effectively sum over thin slabs parallel to the plane of the sky and separated by distances large compared to their depth. One may regard the slab family as mathematically spanning the contributions of the actual physical filaments.

For low galactic latitudes, where most of the neutral hydrogen is located, there may indeed be a preference for filaments alignment in the above thin slab geometry, as a result of the galactic differential rotation. In the SMC the rotation with respect to the Galaxy is expected to have a smaller effect.

In the next section, we show that the observational power spectra are essentially the sum of the spectra contributed by each such thin slab. The mutual contributions coming from *different* slabs are much smaller.

3. H_I Spatial Power Spectra and ISM Turbulence

We wish to relate the observational two-dimensional power spectrum with the underlying three-dimensional turbulence. This will provide an interpretation of the observational power spectrum and at the same time could hopefully yield clues regarding the characteristics of the underlying 3-dimensional turbulence.

The observational 2-dimensional power spectrum is defined as the Fourier transform, over the plane of the sky, of the autocorrelation of the fluctuating brightness temperature $\theta_B = T_B - \langle T_B \rangle$. Here T_B is the brightness temperature and $\langle T_B \rangle$ its average over the plane of the sky. Thus, the power spectrum $p(\vec{q})$ as function of the 2-dimensional wavenumber in the plane of the sky, $\vec{q}$, is

$$p(\vec{q}) = \frac{1}{(2\pi)^2} \int \langle \theta_B(\vec{R}) \theta_B(\vec{R} + \vec{r}) \rangle e^{-i\vec{q}\cdot\vec{r}} d^2r . \quad (2)$$

Here $\vec{R}$, and $\vec{r}$ are in the plane of the sky, and $\langle \theta_B(\vec{R})\theta_B(\vec{R}+\vec{r}) \rangle$ is the autocorrelation of θ_B obtained by averaging over all positions $\vec{R}$ in the plane of the sky. Assuming homogeneity and isotropy in this plane, the power spectrum depends only on the absolute value of the wavenumber, so that $p(\vec{q}) = p(q)$ with $q = |\vec{q}|$. In the optically thin case, the brightness temperature is proportional to the column density of neutral hydrogen in the line of sight (Spitzer 1978), thus in the present case

$$T_B(\vec{R}) = BN_{\text{HI}}(\vec{R}) = B \sum_i \int_{d_i-L_i/2}^{d_i+L_i/2} n(\vec{R}, z) dz \quad (3)$$

with n the number density and B a constant. The summation is over the disjoint thin slabs with depth L_i , located at distances d_i . As mentioned before, each depth L_i is *much smaller* than the slab extent on the plan of the sky and also *much smaller* than the distances between slabs. The fluctuating part of the brightness temperature is thus expressed as

$$\theta_B(\vec{R}) = B \sum_i \int_{d_i-L_i/2}^{d_i+L_i/2} \tilde{n}(\vec{R}, z) dz \quad (4)$$

where $\tilde{n} = n - \langle n \rangle$ is the fluctuating number density, and $\langle n \rangle$ is the average number density in the slab. The autocorrelation of θ_B is thus

$$\langle \theta_B(\vec{R})\theta_B(\vec{R}+\vec{r}) \rangle = B^2 \sum_{i,j} \int_{d_i-L_i/2}^{d_i+L_i/2} \int_{d_j-L_j/2}^{d_j+L_j/2} \langle \tilde{n}(\vec{R}, z')\tilde{n}(\vec{R}+\vec{r}, z) \rangle dz dz' \quad (5)$$

where, again, the angular brackets represent averaging over all positions $\vec{R}$ in the plane of the sky.

From Crovisier & Dickey (1983), Green (1993), and Stanimirović *et al.* (1999) it can be estimated that, typically, the root mean square of θ_B , $\theta_{B\text{rms}} \simeq (0.1 \div 0.2) \langle T_B \rangle$ implying that the root mean square of $\tilde{n}$, $\tilde{n}_{\text{rms}} \simeq (0.1 \div 0.2) \langle n \rangle$. Thus, although the medium is compressible, the density fluctuations giving rise to the brightness temperature fluctuations are not large. This renders plausible the assumption that the density fluctuations are driven by a velocity turbulence and are actually a “passive scalar”, i. e. trace the velocity fluctuations but *do not feed back* upon them. In this case, the 2-point correlation of a density fluctuations field driven by a turbulent velocity field, is proportional to the 2-point correlation of the latter (see e.g. Lesieur 1997)

$$\langle \tilde{n}(\vec{R}, z')\tilde{n}(\vec{R}+\vec{r}, z) \rangle = A^2 \langle \vec{v}(\vec{R}, z') \cdot \vec{v}(\vec{R}+\vec{r}, z) \rangle \quad (6)$$

with A a constant. For simplicity we further assume that the underlying 3-dimensional turbulence is homogeneous and isotropic so that the plane average equals the ensemble average

$$< \vec{v}(\vec{R}, z') \cdot \vec{v}(\vec{R} + \vec{r}, z) > = C_3(\vec{r}, z - z') \quad (7)$$

where C_3 is the 2-point correlation of the underlying 3-dimensional velocity turbulence, defined in terms of *ensemble* average

$$C_3(\vec{r}) = < \vec{v}(\vec{r}') \cdot \vec{v}(\vec{r} + \vec{r}') > = \int \Phi(\vec{k}) e^{i\vec{k} \cdot \vec{r}} d^3k \quad (8)$$

In equation (8), unlike in the previous equations, the position vectors are 3-dimensional and the brackets denote *ensemble* average. C_3 is expressed in terms of the 3-dimensional spectrum $\Phi(\vec{k})$ which in the homogeneous and isotropic case depends on k , the absolute value of the 3-dimensional wavenumber $\vec{k}$,

$$\Phi(\vec{k}) = \Phi(k) = \frac{F(k)}{4\pi k^2} . \quad (9)$$

Here, $F(k)$ is the spectral function of the 3-dimensional velocity turbulence. It equals twice the kinetic energy per unit mass and unit wavenumber in the turbulence.

Accordingly, the mean squared value of the turbulent velocity is

$$v_t^2 = \int_{k_0}^{\infty} F(k) dk \quad (10)$$

with πk_0^{-1} being the largest scale in the turbulence.

We can now go back to equation (2) and use equations (5), (6), (8) and (9) to express $p(q)$ in terms of $F(k)$

$$p(\vec{q}) = \frac{A^2 B^2}{(2\pi)^2} \sum_{i,j} \int_{d_i - L_i/2}^{d_i + L_i/2} \int_{d_j - L_j/2}^{d_j + L_j/2} \int \Phi(\vec{q}', k_z) e^{i\vec{q}' \cdot \vec{r} - i\vec{q} \cdot \vec{r} + i k_z (z - z')} dk_z d^2 q' d^2 r dz dz' , \quad (11)$$

leading to

$$p(\vec{q}) = A^2 B^2 \int \Phi(\vec{q}, k_z) \left(\sum_{i,j} \int_{d_i - L_i/2}^{d_i + L_i/2} \int_{d_j - L_j/2}^{d_j + L_j/2} e^{i k_z (z - z')} dz dz' \right) dk_z = \quad (12)$$

$$A^2 B^2 \int \Phi(\vec{q}, k_z) \left(\sum_i L_i^2 \frac{\sin^2(k_z L_i/2)}{(k_z L_i/2)^2} + \sum_{i \neq j} L_i L_j e^{i k_z (d_i - d_j)} \frac{\sin(k_z L_i/2)}{k_z L_i/2} \frac{\sin(k_z L_j/2)}{k_z L_j/2} \right) dk_z .$$

Since $|d_i - d_j| \gg L_i, L_j$, the contribution of the second term which results from inter-slab correlations is *much smaller* than that of the first term that is due to correlations *within* each slab. Keeping only the first term gives

$$p(\vec{q}) = A^2 B^2 \sum_i L_i^2 \int_0^{\infty} \Phi(\vec{q}, k_z) \frac{\sin^2(k_z L_i/2)}{(k_z L_i/2)^2} dk_z . \quad (13)$$

The assumed homogeneity and isotropy in the plane implies that $p(\vec{q}) = p(q)$, and use of equation (9) leads to

$$p(q) = \frac{A^2 B^2}{4\pi} \sum_i L_i^2 \int_0^\infty \frac{F(\sqrt{q^2 + k_z^2})}{(q^2 + k_z^2)} \frac{\sin^2(k_z L_i/2)}{(k_z L_i/2)^2} dk_z \quad (14)$$

with q being the absolute value of the 2-dimensional in-plane wavenumber, $\vec{q}$. Here k_z denotes the wavenumber in the z direction, perpendicular to the plane of the sky, so that $k^2 = q^2 + k_z^2$. The power spectrum is thus the sum of the power spectra due to each of the thin slabs. Equation (14) determines the power spectrum $p(q)$ for any given spectral function $F(k)$ of the underlying 3-dimensional turbulence .

4. Application to the Observational Power Spectra

It is of interest to use equation (14) in order to deduce information about $F(k)$ from the observational power spectrum $p(q)$. In the general case this cannot be done uniquely. Nevertheless, for power-law functions it is possible.

Since the depth of each slab is much smaller than the observed scales in the plane of the sky for which the power spectra were obtained, the wavenumbers satisfy $q \ll \pi/L_i$. From equation (14) follows that in this case, for an underlying 3-dimensional turbulence spectrum which is a power-law, $F(k) \propto k^{-m}$, the 2-dimensional power spectrum is also a power-law, $p(q) \propto q^{-m-1}$. To see this, note that each of the integrals in equation (14) is now of the form

$$\int_0^\infty (q^2 + k_z^2)^{-m/2-1} \frac{\sin^2(k_z L_i/2)}{(k_z L_i/2)^2} dk_z = I(q) q^{-m-1} \quad (15)$$

where

$$I(q) = \int_0^\infty (1 + y^2)^{-m/2-1} \frac{\sin^2(yqL_i/2)}{(yqL_i/2)^2} dy \quad (16)$$

and $y = k_z/q$.

A plot of $I(q)$ is presented in figure 1 for three representative values of $m = 1, 2, 3$. The wavenumber q is in units of q_0 , corresponding to the largest scale in the plane of the sky. A value $q_0 L_i/\pi = 0.005$ typical to the Galaxy was adopted; for the SMC a typical value is 0.002. For each m , $I(q)$ is essentially constant down to scales in the plane of the sky that are ~ 100 times smaller than the largest scale. Only for very small scales, which turn to be smaller than the observed ones, $I(q) \propto q^{-1}$. For the SMC value of $q_0 L_i/\pi$, the range of scales for which $I(q)$ is constant, would be wider. Thus, each of the terms in equation (14) is a power-law with the *same* index $-(m+1)$.

The above results can be understood qualitatively quite easily. The term $\sin^2(k_z L_i/2)/(k_z L_i/2)^2$ effectively limits the integration over k_z to the range $\sim 0 \div \pi/L_i$. Approximating it by its average value, yields

$$I(q) \propto \int_0^{\pi/(L_i q)} (1 + y^2)^{-m/2-1} dy \quad (17)$$

Indeed, in the limit of $q \ll \pi/L_i$ which applies here, $I(q)$ is constant.

The power spectrum for each of the velocity groups is a sum of the spectra of the individual slabs. Since each of these power spectra is a power-law with the *same* index, the combined power spectrum will also be a power-law with this index. We conclude that a power spectrum $p(q) \propto q^{-m-1}$ is the result of an underlying 3-dimensional turbulence characterized by a spectral function $F(k) \propto k^{-m}$. The observational values of the index of the power spectrum, imply that $m \sim 1.8 \div 2$ for the Galaxy and $m \sim 2$ for the SMC.




Fig. 1.— $\text{Log } I(q)$ as function of $\text{Log } q$, (in units of q_0), for $q_0 L_i / \pi = 0.005$. The curves correspond to $m = 1, 2, 3$ in this order, with $m = 1$ being the uppermost one.

5. Discussion

To interpret the observational spatial power spectra we have assumed that the density fluctuations that produce them are a response to a large scale 3-dimensional turbulence in the ISM, in which the emitting neutral hydrogen regions are embedded.

We derived an expression for the intensity power spectrum in terms of the underlying 3-dimensional large scale turbulence.

The power spectra were seen to be the sum of contributions from thin slabs of neutral hydrogen, parallel to the plane of the sky, with depth *much smaller* than their extent in the plane of the sky. The depth is also *much smaller* than the distances, along the line of sight, between slabs. In this case, we found that a power-law 3-dimensional turbulence spectrum, $F(k) \propto k^{-m}$, leads to a power-law 2-dimensional intensity power spectrum, $p(q) \propto q^{-m-1}$. From the observational values of the power spectrum indices, it follows that $m \sim 2$ for the SMC and $m = 1.8 - 2$ for the Galaxy.

5.1. Turbulence spectrum

This 3-dimensional turbulence spectrum resembles the Kolmogorov spectrum characterized by $m = 5/3$ but differs in the value of the index which is closer to $m = 2$. For the Kolmogorov spectrum the turbulent velocity on scale l satisfies $v_t(l) \propto l^{1/3}$, while in the present case $v_t(l) \propto l^{1/2}$. A velocity-size relation with index ~ 0.5 , has been deduced for molecular clouds of a wide range of sizes (Larson 1981; Leung, Kutner & Mead 1982; Myers 1983; Dame *et al.* 1986; Myers & Goodman 1988; Falgarone, Puget & Perault 1992; Miesch & Bally 1994). A turbulence spectrum characterized by $m = 2$ was also found in a H_{II} region by Roy & Joncas (1985). It is interesting that this spectrum seems to prevail in different components of the ISM. The common factor in these cases is that the medium is compressible. Indeed, numerical simulations, with and without a magnetic field (Passot, Pouquet, & Woodward 1988, Vázquez-Samedani, Ballesteros-Paredes, & Rodríguez 1997), indicate that such a spectrum is characteristic to turbulence in a compressible medium (whether or not magnetic field fluctuations are important).

The Kolmogorov spectrum strictly applies to the inertial wavenumber range for homogeneous and isotropic incompressible turbulence. In this range there is no energy input from the source generating the turbulence (input is at larger scales) nor energy losses (dissipation is at smaller scales). Thus, turbulent kinetic energy is cascaded from larger to smaller scales due to the nonlinear eddy interactions, at a *constant rate* that is *independent of wavenumber*.

The present steeper spectrum, with $m \sim 2$, indicates that at each wavenumber a part of the cascaded turbulent kinetic energy is transformed to other energy forms. As a result, the turbulent kinetic energy rate is smaller the larger the wavenumber. In a compressible medium, part of the kinetic turbulent energy is indeed transformed to energy associated with density fluctuations.

Finally, we note that the emitting neutral regions themselves, modeled here as thin slabs, are probably transient features created and disrupted by the large scale turbulence (Ballesteros-Paredes, Vázquez-Samedani, & Scalo 1999). Their life-times

are of the order of the timescales of the largest eddies. The spatial scales for which the power spectra were obtained are smaller than the above, and therefore the corresponding turbulence timescales are shorter than those of the very largest scales. With respect to the observed scales, the slab can be considered as a quasi-stationary structure in which the observed smaller scale eddies generate the density fluctuations that give rise to the intensity fluctuations.

5.2. Turbulence Scales and Energy Sources

The scale of the large-scale turbulence in the Galaxy should exceed 200 pc – the largest scale probed by the power spectra. The power-law shape of the spectra suggests that energy input to the turbulence is on yet larger scales. A probable energy source on these larger scales is the galactic differential rotation, as also suggested by numerical simulations of Wada & Norman (1999). In this case, a natural scale will be a fraction of the galactocentric distance. However, for a 3- dimensional turbulence, the largest scale is likely to be comparable to the width of the galactic H_{II} layer of ~ 2 kpc (Reynolds 1989; Reynolds 1991).

In the SMC, the turbulence can be generated by a host of instabilities related to the bulk flows that result from the tidal interaction with the Galaxy and with the LMC, as well as to the expanding shells and super shells. The existence of large scale correlated density fluctuations in the SMC indicates the existence of turbulence on a scale $\gtrsim 4$ kpc which is comparable to the size of the SMC . This observational evidence by itself is a very interesting result as the turbulence scale is an order of magnitude larger than that of the turbulence in the LMC, studied by Spicker & Feitzinger (1988) on the basis of radial velocities of 21-cm emission.

The most plausible source feeding energy to such an extremely large scale turbulence are bulk flows induced by the tidal interactions with the Galaxy and the LMC, notably the very close encounter with the LMC that peaked about 2×10^8 yr ago and lasted for about 2×10^8 yr (Gardiner & Noguchi 1996). Such large scale bulk flows generate turbulence due to shear instabilities e. g. the Kelvin-Helmholtz instability (KH) or shock induced Richtmier- Meshkov instability(RM). These two mechanisms were shown to induce large scale turbulence in the intracluster gas of merging galaxy clusters by Roettiger *et al.* (1998) and by Goldman (1997), respectively.

The simulations of Gardiner & Noguchi (1996) exhibit large scale (few kpc) velocity gradients with shear values $S \simeq 5 - 10$ km/s/kpc, consistent with the (long period, population I) Cepheid radial velocity data of Mathewson, Ford,& Visvanathan (1986, 1988). One expects that the large scale shear will produce turbulence with the largest eddies size l_t comparable to these spatial scales, and root mean squared turbulent velocities $v_t \lesssim Sl_t = 10 \div 15$ km/s. This value is comparable to the sound speed of the embedding ionized medium, and supersonic with respect to the H_I regions, thus

enabling the density fluctuations in the latter. On smaller scales of $\lesssim 1$ kpc, also the giant super-shells observed in the SMC, contribute, to the turbulence. The giant super-shells themselves are probably the result of a star formation burst that followed the interaction with the LMC (Zaritsky *et al.* 2000).

The time scale for decay of the turbulence $t_d \sim l_t/v_t$ with a rms turbulent velocity of $v_t \approx 10$ km/s and scale $l_t = 1$ kpc is $\sim 4 \times 10^8$ yr. Thus, we are witnessing a turbulence that didn't have enough time to decay, since the close encounter with the LMC of $\sim 2 \times 10^8$ yr ago. Moreover, the simulations of Roettiger *et al.* (1998) and simulations of RM instability (Rotman, 1991) indicate that continued excitation (even at a lower level than the original one) can keep such a turbulence going for times considerably longer than the above estimate. Such a lower level excitation can be provided by ongoing tidal interactions of smaller' strength.

Another question of interest is the near constancy of the slope of the spectra. We know that small scale (from 1 kpc and down)) energy sources exist (supergiant shells, smaller scale supernovae remnants, smaller scale hydrodynamic instabilities) as well as smaller scale energy sinks (e.g. magnetic excitations). Consider a source /sink at wavenumber k in an interval Δk such that $k \gtrsim \Delta k \gg k_0$ with k_0 corresponding to the largest scale in the turbulence. Denote by ϵ and $\Delta\epsilon$ the energy rate due to the large scale turbulence and the change due to the small scale source, respectively. Since $F(k) \propto \epsilon(k)^{2/3}$ the jump in the turbulence spectral function

$$\frac{\Delta F(k)}{F(k)} \sim \frac{2}{3} \frac{\Delta\epsilon}{\epsilon} . \quad (18)$$

Even for a relative change of $\sim 20\%$, the jump will be $\lesssim 10\%$, consistent with the scatter of the observational spectra.

5.3. Implications of the Turbulence on the SMC Magnetic Field and Abundances Gradients

The consequences of this turbulence may be quite important regarding magnification of large scale magnetic fields, and enhancing diffusion coefficients. In the SMC, the turbulence largest eddies are comparable in scale to the size of the SMC. This implies that turbulent mixing should smooth any large scale abundance gradients. This seems indeed to be the case observationally (Dufour 1975; Kobulnicky 1998) . The turbulence is also expected to amplify and shape up the large scale magnetic field. Indeed, the SMC observational data indicate the existence of a large scale disordered field (Haynes *et al.* 1991; Ye & Turtle 1991). The observed magnetic field is $\sim 6.5 \mu\text{G}$, a value close to that expected from a dynamo amplification to an equilibrium value $B = 5 \mu\text{G} (v_t/10 \text{ km s}^{-1}) (n/1 \text{ cm}^{-3})^{1/2}$ with v_t the turbulent root mean squared velocity and n the number density in the ionized embedding region.

5.4. Concluding Remarks

The interpretation of the H_I power spectrum of the SMC in terms of an underlying large scale turbulence seems natural and as noted above is consistent with other observational data. The existence of a large scale turbulence in the SMC gains also support from the recent finding of Stanimirović *et al.* (2000), that the autocorrelation of the dust column density is also a power law with an index ~ -3 . In our interpretation, the dust will be just another passive marker waved by the turbulence and thus should have the same power-law spectral index as the H_I intensity.

Moreover, observations of stars of various ages were used by Zaritsky *et al.* (2000) and by Kunkel, Demers, & Irwin (2000), to conclude that hydrodynamic interactions, following the recent close passage of the LMC, rather than just gravity are responsible for the morphology and dynamics of the ISM of the SMC.

Comments by the referee and by Dr. Steven Shore have contributed to the improvement of the presentation. This work has been supported by the US-Israel BSF grant 94-314 and Israel Science Foundation grant 561-9911.2.

REFERENCES

- Ballesteros-Paredes, J., Vázquez-Samedani, E., & Scalo, J. 1999, ApJ, 515, 286
- Crovisier, J., & Dickey, J. M. 1983, A&A, 122, 282
- Dame, T. M., Elemgreen, B. G., Cohen, R. S., Thaddeus, P. 1986, ApJ, 305, 892
- Dufour, R. J. 1975, ApJ, 195, 315
- Falgarone, E., Puget, J.-L., & Perault, M. 1992, A&A, 257, 715
- Gardiner, L. T., & Noguchi, M. 1996, MNRAS, 278,191
- Goldman, I. 1998, In Proceedings of Iau symposium 188 "The Hot Universe", Kyoto, Japan The hot universe edited by Katsuji Koyama; Shunji Kitamoto; Masayuki Itoh. Dordrecht : Kluwer Sym. No.188., p.297
- Green, D. A. 1993, MNRAS, 262, 327
- Haynes, R. F., Klein, U., Wayte, S.R., Wielebinski, R, Murray, J. D., Bajaaaaaja, E., Meinert, D., Buczilowski, Harnett, J.I., Hunt, A.J., Wark, R., & Sciacca, L. 1991, A&A, 252, 475
- Kobulnicky, C. 1998, PASP Conference Series, 147,108
- Kunkel, W. E., Demers, S., & Irwin, M. J. 2000, astro-ph/0003395
- Lazarian, A. 1995, A&A, 293, 507
- Lazarian, A. & Pogosayn, D. 2000, ApJ, accepted
- Larson, R. B. 1981, MNRAS,194, 809

- Lesiurs, M. 1997, *Turbulence in Fluids*, §6.10 (Dordrecht: Kluwer)
- Leung, C. M., Kutner, M. L., & Mead, K. N. 1982, *ApJ*, 262, 583
- Mathewson, D. S., Ford, V. L., & Visvanathan, N. 1986 *ApJ*, 301, 664
- Mathewson, D. S., Ford, V. L., & Visvanathan, N. 1988 *ApJ*, 333, 617
- Miesch, M. S., & Bally, J. 1994, *ApJ*, 429, 645
- Myers, P. C. 1983, *ApJ*, 270, 105
- Myers, P. C., & Goodman, A. A. 1988, *ApJ*, 329, 392
- Passot, T., Pouquet, A., & Woodward, P. 1988, *A&A*, 197, 228
- Reynolds, R. J. 1989, *ApJ*, 339, L29
- Reynolds, R. J. 1991, in *The Interstellar Disk-Halo Connection in Galaxies*, IAU Symp. No.144, ed. H. Bloemen (Dordrecht: Kluwer), p. 67
- Roettiger, K., Stone, J. M., & Mushotzky, R. F. 1998, *ApJ*, 493, 62
- Rotman, D. 1991, *Phys. Fl.* A3, 1792;
- Roy, J. R., & Joncas, G. 1985, *ApJ*, 288, 142
- Spicker, J., & Feitzinger, J. V. 1988, *A&A*, 191, 10
- Spitzer, L. Jr. 1978, *Physical Processes in the Interstellar Medium* (New York:Wiley)
- Stanimirović, S., Stavely-Smith, L., Dickey, J. M., Sault, R. J., & Snowden, S. L. 1999, *MNRAS*, 302, 417
- Stanimirović, S., Stavely-Smith, L., van der Hulst, J. M., Bontekoe, Tj. R., Kester, D. J. M., & Jones, P. A. 2000, *MNRAS*
- Ye, T., & Turtle, A.J., 1991, *MNRAS*, 249, 693
- Vázquez-Samedani, E., Ballesteros-Paredes, J., & Rodríguez, L. F. 1997, *ApJ*, 474, 292
- Wada, K., & Norman, C. A. 1999, *ApJ*, 516, L13
- Wolfire, M. G., Hollenbach, D., McKee, C. F., Tielens, A. G. G. M., & Bakes, E. L. O. 1995, *ApJ*, 443, 152
- Zaritsky, D., Harris, J., Grebel, E. K., & Thompson, I. B. 2000, *astro-ph/0003155*